

Initial test analyte	Initial test cutoff concentration	Confirmatory test analyte	Confirmatory test cutoff concentration
Marijuana metabolites	50 ng/ml	THCA ¹	15ng/ml
Cocaine metabolites	150 ng/ml	Benzoylcegonine	100 ng/ml
Opiate metabolites. Codeine/Morphine ²	2000 ng/ml	Codeine Morphine	2000 ng/ml 2000 ng/ml
6-Acetylmorphine ³	10 ng/ml	6-Acetylmorphine	10 ng/ml
Phencyclidine	25 ng/ml	Phencyclidine	25 ng/ml
Amphetamines. AMP/MAMP ⁴	500 ng/ml	Amphetamine Methamphetamine ⁵	250 ng/ml 250 ng/ml
MDMA ⁶	500 ng/ml	MDMA MDA ⁷ MDEA ⁸	250 ng/ml 250 ng/ml 250 ng/ml

¹ Delta-9-tetrahydrocannabinol-9-carboxylic acid (THCA).

² Morphine is the target analyte for codeine/morphine testing.

³ Either a single initial test kit or multiple initial test kits may be used provided the single test kit detects each target analyte independently at the specified cutoff.

⁴ Methamphetamine is the target analyte for amphetamine/methamphetamine testing.

⁵ To be reported positive for methamphetamine, a specimen must also contain amphetamine at a concentration equal to or great than 100 ng/ml.

⁶ Methylenedioxyamphetamine (MDMA).

⁷ Methylenedioxyamphetamine (MDA).